

PATENT  
Docket: 014/002C

region of the hTR, wherein the first nucleotide sequence within the accessible region is selected from the group consisting of nucleotides 137-196, nucleotides 290-319, and nucleotides 350-380 of hTR, and wherein the polynucleotide comprises a nucleotide analog or a non-naturally occurring nucleotide linkage selected from the group consisting of phosphorothioates, phosphoramidates, methyl phosphonates, chiral-methyl phosphonates, 2-O-methyl ribonucleotides and peptide-nucleic acids.

28. (new) A polynucleotide according to claim 27, when the polynucleotide sequence is selected from the group consisting of:

CGT TCC TCT TCC TGC GGC CTG AAA CGG TGA (SEQ ID NO:2)  
CGT TCC TCT TCC TGC GGC CT (SEQ ID NO:3)  
CGT TCC TCT TCC (SEQ ID NO:4)  
CTG ACA GAG CCC AAC TCT TCG CGG TGG CAG (SEQ ID NO. 5)  
CTG ACA GAG CCC AAC TCT TC (SEQ ID NO:6)  
CCA ACT CTT CGC GGT GGC AG (SEQ ID NO:7)  
GCT CTA GAA TGA ACG GTG GAA GGC GGC AGG (SEQ ID NO:8)  
GCT CTA GAA TGA ACG GTG G (SEQ ID NO.9)  
GCT CTA GAA TGA ACG (SEQ ID NO:10)  
GCT CTA GAA TG (SEQ ID NO:11)  
GCT CTA G (SEQ ID NO:12)  
CAT TTT TTG TTT GCT CTA GA (SEQ ID NO:13) and  
CGG GCC AGC AGC TGA CA (SEQ ID NO:14).

29. (new) A polynucleotide consisting essentially of a sequence selected from the group consisting of:

CGT TCC TCT TCC TGC GGC CTG AAA CGG TGA (SEQ ID NO:2)  
CGT TCC TCT TCC TGC GGC CT (SEQ ID NO:3)  
CGT TCC TCT TCC (SEQ ID NO:4)  
CTG ACA GAG CCC AAC TCT TCG CGG TGG CAG (SEQ ID NO. 5)  
CTG ACA GAG CCC AAC TCT TC (SEQ ID NO:6)  
CCA ACT CTT CGC GGT GGC AG (SEQ ID NO:7)